Journal of Integer Sequences, Vol. 10 (2007),
Article 07.8.2

The Compositions of Differential Operations and the Gateaux Directional Derivative

Branko J. Malešević[1] and Ivana V. Jovović[2]
University of Belgrade
Faculty of Electrical Engineering
Bulevar kralja Aleksandra 73
Belgrade
Serbia
malesh@EUnet.yu
ivana121@EUnet.yu

Abstract

This paper deals with the number of meaningful compositions of higher order of differential operations and the Gateaux directional derivative.

1 The compositions of differential operations of the space $\mathbb{R}^3$

In the real three-dimensional space $\mathbb{R}^3$ we consider the following sets:

$$\mathrm{A}_0 = \{f : \mathbb{R}^3 \longrightarrow \mathbb{R} \mid f \in C^\infty(\mathbb{R}^3)\} \quad \text{and} \quad \mathrm{A}_1 = \{\vec{f} : \mathbb{R}^3 \longrightarrow \mathbb{R}^3 \mid \vec{f} \in \vec{C}^\infty(\mathbb{R}^3)\}. \tag{1}$$

It is customary in vector analysis to consider $m = 3$ basic differential operations on A_0 and A_1 [1], namely:

$$\begin{aligned}
&\operatorname{grad} f = \nabla_1 f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \frac{\partial f}{\partial x_3}\right) : \mathrm{A}_0 \longrightarrow \mathrm{A}_1, \\
&\operatorname{curl} \vec{f} = \nabla_2 \vec{f} = \left(\frac{\partial f_3}{\partial x_2} - \frac{\partial f_2}{\partial x_3}, \frac{\partial f_1}{\partial x_3} - \frac{\partial f_3}{\partial x_1}, \frac{\partial f_2}{\partial x_1} - \frac{\partial f_1}{\partial x_2}\right) : \mathrm{A}_1 \longrightarrow \mathrm{A}_1, \\
&\operatorname{div} \vec{f} = \nabla_3 \vec{f} = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} + \frac{\partial f_3}{\partial x_3} : \mathrm{A}_1 \longrightarrow \mathrm{A}_0 .
\end{aligned} \tag{2}$$

[1]Supported in part by the project MNTRS, Grant No. ON144020.
[2]Ph. D. student, Faculty of Mathematics, University of Belgrade, Serbia.

Let us present the number of meaningful compositions of higher order over the set $\mathcal{A}_3 = \{\nabla_1, \nabla_2, \nabla_3\}$. It is familiar fact that there are $m = 5$ compositions of the second order [2, p. 161]:

$$\begin{aligned}
&\Delta f = \operatorname{div}\operatorname{grad} f = \nabla_3 \circ \nabla_1 f,\\
&\operatorname{curl}\operatorname{curl} \vec{f} = \nabla_2 \circ \nabla_2 \vec{f},\\
&\operatorname{grad}\operatorname{div} \vec{f} = \nabla_1 \circ \nabla_3 \vec{f},\\
&\operatorname{curl}\operatorname{grad} f = \nabla_2 \circ \nabla_1 f = \vec{0},\\
&\operatorname{div}\operatorname{curl} \vec{f} = \nabla_3 \circ \nabla_2 \vec{f} = 0.
\end{aligned} \tag{3}$$

Malešević [3] proved that there are $m = 8$ compositions of the third order:

$$\begin{aligned}
&\operatorname{grad}\operatorname{div}\operatorname{grad} f = \nabla_1 \circ \nabla_3 \circ \nabla_1 f,\\
&\operatorname{curl}\operatorname{curl}\operatorname{curl} \vec{f} = \nabla_2 \circ \nabla_2 \circ \nabla_2 \vec{f},\\
&\operatorname{div}\operatorname{grad}\operatorname{div} \vec{f} = \nabla_3 \circ \nabla_1 \circ \nabla_3 \vec{f},\\
&\operatorname{curl}\operatorname{curl}\operatorname{grad} f = \nabla_2 \circ \nabla_2 \circ \nabla_1 f = \vec{0},\\
&\operatorname{div}\operatorname{curl}\operatorname{grad} f = \nabla_3 \circ \nabla_2 \circ \nabla_1 f = 0,\\
&\operatorname{div}\operatorname{curl}\operatorname{curl} \vec{f} = \nabla_3 \circ \nabla_2 \circ \nabla_2 \vec{f} = 0,\\
&\operatorname{grad}\operatorname{div}\operatorname{curl} \vec{f} = \nabla_1 \circ \nabla_3 \circ \nabla_2 \vec{f} = \vec{0},\\
&\operatorname{curl}\operatorname{grad}\operatorname{div} \vec{f} = \nabla_2 \circ \nabla_1 \circ \nabla_3 \vec{f} = \vec{0}.
\end{aligned} \tag{4}$$

If $\mathtt{f}(k)$ is the number of compositions of the kth order, then Malešević [4] proved

$$\mathtt{f}(k) = F_{k+3}, \tag{5}$$

where F_k is k^{th} Fibonacci number.

2 The compositions of the differential operations and Gateaux directional derivative of the space $\mathbb{R}^3$

Let $f \in \mathrm{A}_0$ be a scalar function and $\vec{e} = (e_1, e_2, e_3) \in \mathbb{R}^3$ be a unit vector. The *Gateaux directional derivative* in direction $\vec{e}$ is defined by [5, p. 71]:

$$\operatorname{dir}_{\vec{e}} f = \nabla_0 f = \nabla_1 f \cdot \vec{e} = \frac{\partial f}{\partial x_1} e_1 + \frac{\partial f}{\partial x_2} e_2 + \frac{\partial f}{\partial x_3} e_3 : \mathrm{A}_0 \longrightarrow \mathrm{A}_0. \tag{6}$$

Let us determine the number of meaningful compositions of higher order over the set $\mathcal{B}_3 = \{\nabla_0, \nabla_1, \nabla_2, \nabla_3\}$. There exist $m = 8$ compositions of the second order:

$$
\begin{aligned}
&\operatorname{dir}_{\vec{e}} \operatorname{dir}_{\vec{e}} f = \nabla_0 \circ \nabla_0 f = \nabla_1\big(\nabla_1 f \cdot \vec{e}\big) \cdot \vec{e},\\
&\operatorname{grad} \operatorname{dir}_{\vec{e}} f = \nabla_1 \circ \nabla_0 f = \nabla_1\big(\nabla_1 f \cdot \vec{e}\big),\\
&\Delta f = \operatorname{div} \operatorname{grad} f = \nabla_3 \circ \nabla_1 f,\\
&\operatorname{curl} \operatorname{curl} \vec{f} = \nabla_2 \circ \nabla_2 \vec{f},\\
&\operatorname{dir}_{\vec{e}} \operatorname{div} \vec{f} = \nabla_0 \circ \nabla_3 \vec{f} = \big(\nabla_1 \circ \nabla_3 \vec{f}\big) \cdot \vec{e},\\
&\operatorname{grad} \operatorname{div} \vec{f} = \nabla_1 \circ \nabla_3 \vec{f},\\
&\operatorname{curl} \operatorname{grad} f = \nabla_2 \circ \nabla_1 f = \vec{0},\\
&\operatorname{div} \operatorname{curl} \vec{f} = \nabla_3 \circ \nabla_2 \vec{f} = 0;
\end{aligned}
\tag{7}
$$

and there exist $m = 16$ compositions of the third order:

$$
\begin{aligned}
&\operatorname{dir}_{\vec{e}} \operatorname{dir}_{\vec{e}} \operatorname{dir}_{\vec{e}} f = \nabla_0 \circ \nabla_0 \circ \nabla_0 f,\\
&\operatorname{grad} \operatorname{dir}_{\vec{e}} \operatorname{dir}_{\vec{e}} f = \nabla_1 \circ \nabla_0 \circ \nabla_0 f,\\
&\operatorname{div} \operatorname{grad} \operatorname{dir}_{\vec{e}} f = \nabla_3 \circ \nabla_1 \circ \nabla_0 f,\\
&\operatorname{dir}_{\vec{e}} \operatorname{div} \operatorname{grad} f = \nabla_0 \circ \nabla_3 \circ \nabla_1 f,\\
&\operatorname{grad} \operatorname{div} \operatorname{grad} f = \nabla_1 \circ \nabla_3 \circ \nabla_1 f,\\
&\operatorname{curl} \operatorname{curl} \operatorname{curl} \vec{f} = \nabla_2 \circ \nabla_2 \circ \nabla_2 \vec{f},\\
&\operatorname{dir}_{\vec{e}} \operatorname{dir}_{\vec{e}} \operatorname{div} \vec{f} = \nabla_0 \circ \nabla_0 \circ \nabla_3 \vec{f},\\
&\operatorname{grad} \operatorname{dir}_{\vec{e}} \operatorname{div} \vec{f} = \nabla_1 \circ \nabla_0 \circ \nabla_3 \vec{f},\\
&\operatorname{div} \operatorname{grad} \operatorname{div} \vec{f} = \nabla_3 \circ \nabla_1 \circ \nabla_3 \vec{f},\\
&\operatorname{curl} \operatorname{grad} \operatorname{dir}_{\vec{e}} f = \nabla_2 \circ \nabla_1 \circ \nabla_0 \vec{f} = \vec{0},\\
&\operatorname{curl} \operatorname{curl} \operatorname{grad} f = \nabla_2 \circ \nabla_2 \circ \nabla_1 f = \vec{0},\\
&\operatorname{div} \operatorname{curl} \operatorname{grad} f = \nabla_3 \circ \nabla_2 \circ \nabla_1 f = 0,\\
&\operatorname{div} \operatorname{curl} \operatorname{curl} \vec{f} = \nabla_3 \circ \nabla_2 \circ \nabla_2 \vec{f} = 0,\\
&\operatorname{dir}_{\vec{e}} \operatorname{div} \operatorname{curl} \vec{f} = \nabla_0 \circ \nabla_3 \circ \nabla_2 \vec{f} = 0,\\
&\operatorname{grad} \operatorname{div} \operatorname{curl} \vec{f} = \nabla_1 \circ \nabla_3 \circ \nabla_2 \vec{f} = \vec{0},\\
&\operatorname{curl} \operatorname{grad} \operatorname{div} \vec{f} = \nabla_2 \circ \nabla_1 \circ \nabla_3 \vec{f} = \vec{0}.
\end{aligned}
\tag{8}
$$

Further on we shall use the method from the paper [4]. Let us define a binary relation σ "*to be in composition*": $\nabla_i \, \sigma \, \nabla_j$ iff the composition $\nabla_j \circ \nabla_i$ is meaningful. Then Cayley table of the relation σ is determined by

σ	∇_0	∇_1	∇_2	∇_3
∇_0	$\top$	$\top$	$\bot$	$\bot$
∇_1	$\bot$	$\bot$	$\top$	$\top$
∇_2	$\bot$	$\bot$	$\top$	$\top$
∇_3	$\top$	$\top$	$\bot$	$\bot$

(9)

Let us denote by ∇_{-1} nowhere-defined function, where domain and range are empty sets [3] and let $\nabla_{-1} \, \sigma \, \nabla_i$ hold for $i = 0, 1, 2, 3$. If G is graph which is determined by the relation σ, then graph of paths of G is the tree with the root ∇_{-1} (Fig. 1).



Fig. 1

Let $\mathsf{g}(k)$ be the number of meaningful compositions of the kth order of the functions from $\mathcal{B}_3$ and let $\mathsf{g}_i(k)$ be the number of meaningful compositions of the kth order beginning from the left by ∇_i. Then $\mathsf{g}(k) = \mathsf{g}_0(k) + \mathsf{g}_1(k) + \mathsf{g}_2(k) + \mathsf{g}_3(k)$. Based on the partial self similarity of the tree (Fig. 1) we obtain equalities

$$\begin{aligned} \mathsf{g}_0(k) &= \mathsf{g}_0(k-1) + \mathsf{g}_1(k-1), \\ \mathsf{g}_1(k) &= \mathsf{g}_2(k-1) + \mathsf{g}_3(k-1), \\ \mathsf{g}_2(k) &= \mathsf{g}_2(k-1) + \mathsf{g}_3(k-1), \\ \mathsf{g}_3(k) &= \mathsf{g}_0(k-1) + \mathsf{g}_1(k-1). \end{aligned} \tag{10}$$

Hence, the recurrence for $\mathsf{g}(k)$ is

$$\mathsf{g}(k) = 2\,\mathsf{g}(k-1) \tag{11}$$

and because $\mathsf{g}(1) = 4$ we have

$$\mathsf{g}(k) = 2^{k+1}. \tag{12}$$

3 The compositions of differential operations of the space $\mathbb{R}^{\mathbf{n}}$

Let us present the number of meaningful compositions of differential operations in the vector analysis of the space $\mathbb{R}^n$, where differential operations ∇_r $(r = 1, \ldots, n)$ are defined on

corresponding non-empty sets A_s ($s=1,\ldots,m$ and $m=\lfloor n/2\rfloor$, $n\geq 3$) according to the papers [4], [6]:

$$\begin{array}{ll}
\mathcal{A}_n\ (n{=}2m): & \nabla_1:\mathrm{A}_0\to\mathrm{A}_1 \\
& \nabla_2:\mathrm{A}_1\to\mathrm{A}_2 \\
& \vdots \\
& \nabla_i:\mathrm{A}_{i-1}\to\mathrm{A}_i \\
& \vdots \\
& \nabla_m:\mathrm{A}_{m-1}\to\mathrm{A}_m \\
& \nabla_{m+1}:\mathrm{A}_m\to\mathrm{A}_{m-1} \\
& \vdots \\
& \nabla_{n-j}:\mathrm{A}_{j+1}\to\mathrm{A}_j \\
& \vdots \\
& \nabla_{n-1}:\mathrm{A}_2\to\mathrm{A}_1 \\
& \nabla_n:\mathrm{A}_1\to\mathrm{A}_0,
\end{array}
\qquad
\begin{array}{ll}
\mathcal{A}_n\ (n{=}2m{+}1): & \nabla_1:\mathrm{A}_0\to\mathrm{A}_1 \\
& \nabla_2:\mathrm{A}_1\to\mathrm{A}_2 \\
& \vdots \\
& \nabla_i:\mathrm{A}_{i-1}\to\mathrm{A}_i \\
& \vdots \\
& \nabla_m:\mathrm{A}_{m-1}\to\mathrm{A}_m \\
& \nabla_{m+1}:\mathrm{A}_m\to\mathrm{A}_m \\
& \nabla_{m+2}:\mathrm{A}_m\to\mathrm{A}_{m-1} \\
& \vdots \\
& \nabla_{n-j}:\mathrm{A}_{j+1}\to\mathrm{A}_j \\
& \vdots \\
& \nabla_{n-1}:\mathrm{A}_2\to\mathrm{A}_1 \\
& \nabla_n:\mathrm{A}_1\to\mathrm{A}_0.
\end{array}
\tag{13}$$

Let us define *higher order differential operations* as meaningful compositions of higher order of differential operations from the set $\mathcal{A}_n=\{\nabla_1,\ldots,\nabla_n\}$. The number of higher order differential operations is given according to the paper [4]. Furthermore, let us define a binary relation ρ "*to be in composition*": $\nabla_i\,\rho\,\nabla_j$ iff the composition $\nabla_j\circ\nabla_i$ is meaningful. Then Cayley table of the relation ρ is determined by

$$\nabla_i\,\rho\,\nabla_j=\begin{cases}\top & ,\ (j=i+1)\vee(i+j=n+1);\\ \bperp & ,\ \text{otherwise.}\end{cases}\tag{14}$$

Let $\mathtt{A}=[a_{ij}]\in\{0,1\}^{n\times n}$ be the adjacency matrix associated with the graph which is determined by the relation ρ. Malešević [6] proved the following statements.

Theorem 3.1. *Let $P_n(\lambda)=|\mathtt{A}-\lambda\mathtt{I}|=\alpha_0\lambda^n+\alpha_1\lambda^{n-1}+\cdots+\alpha_n$ be the characteristic polynomial of the matrix $\mathtt{A}$ and $v_n=[\,1\ \ldots\ 1\,]_{1\times n}$. If $\mathtt{f}(k)$ is the number of the kth order differential operations, then the following formulas hold:*

$$\mathtt{f}(k)=v_n\cdot\mathtt{A}^{k-1}\cdot v_n^T\tag{15}$$

and

$$\alpha_0\mathtt{f}(k)+\alpha_1\mathtt{f}(k-1)+\cdots+\alpha_n\mathtt{f}(k-n)=0\quad(k>n).\tag{16}$$

Lemma 3.2. *Let $P_n(\lambda)$ be the characteristic polynomial of the matrix* $\mathtt{A}$*. Then the following recurrence holds:*

$$P_n(\lambda) = \lambda^2\big(P_{n-2}(\lambda) - P_{n-4}(\lambda)\big). \tag{17}$$

Lemma 3.3. *Let $P_n(\lambda)$ be the characteristic polynomial of the matrix* $\mathtt{A}$*. Then it has the following explicit form:*

$$P_n(\lambda) = \begin{cases} \displaystyle\sum_{k=1}^{\lfloor\frac{n+2}{4}\rfloor+1} (-1)^{k-1}\binom{\frac{n}{2}-k+2}{k-1}\lambda^{n-2k+2} & , \ n=2m; \\ \displaystyle\sum_{k=1}^{\lfloor\frac{n+2}{4}\rfloor+2} (-1)^{k-1}\left(\binom{\frac{n+3}{2}-k}{k-1}+\binom{\frac{n+3}{2}-k}{k-2}\lambda\right)\lambda^{n-2k+2} & , \ n=2m+1. \end{cases} \tag{18}$$

From previous statements one can obtain the recurrences in the table, [4]:

Dimension	Recurrence for the number of the kth order differential operations
$n = 3$	$\mathtt{f}(k) = \mathtt{f}(k-1) + \mathtt{f}(k-2)$
$n = 4$	$\mathtt{f}(k) = 2\mathtt{f}(k-2)$
$n = 5$	$\mathtt{f}(k) = \mathtt{f}(k-1) + 2\mathtt{f}(k-2) - \mathtt{f}(k-3)$
$n = 6$	$\mathtt{f}(k) = 3\mathtt{f}(k-2) - \mathtt{f}(k-4)$
$n = 7$	$\mathtt{f}(k) = \mathtt{f}(k-1) + 3\mathtt{f}(k-2) - 2\mathtt{f}(k-3) - \mathtt{f}(k-4)$
$n = 8$	$\mathtt{f}(k) = 4\mathtt{f}(k-2) - 3\mathtt{f}(k-4)$
$n = 9$	$\mathtt{f}(k) = \mathtt{f}(k-1) + 4\mathtt{f}(k-2) - 3\mathtt{f}(k-3) - 3\mathtt{f}(k-4) + \mathtt{f}(k-5)$
$n = 10$	$\mathtt{f}(k) = 5\mathtt{f}(k-2) - 6\mathtt{f}(k-4) + \mathtt{f}(k-6)$

The values of the function $\mathtt{f}(k)$, for small values of the argument k, are given in the database of integer sequences [8] as the following sequences A020701 ($n = 3$), A090989 ($n = 4$), A090990 ($n = 5$), A090991 ($n = 6$), A090992 ($n = 7$), A090993 ($n = 8$), A090994 ($n = 9$), A090995 ($n = 10$).

4 The compositions of differential operations and Gateaux directional derivative of the space $\mathbb{R}^n$

Let $f \in \mathrm{A}_0$ be a scalar function and $\vec{e} = (e_1, \ldots, e_n) \in \mathbb{R}^n$ be a unit vector. The *Gateaux directional derivative* in direction $\vec{e}$ is defined by [5, p. 71]:

$$\mathrm{dir}_{\vec{e}}\, f = \nabla_0 f = \sum_{k=1}^{n} \frac{\partial f}{\partial x_k}\, e_k : \mathrm{A}_0 \longrightarrow \mathrm{A}_0. \tag{19}$$

Let us extend the set of differential operations $\mathcal{A}_n = \{\nabla_1, \dots, \nabla_n\}$ with Gateaux directional derivative to the set $\mathcal{B}_n = \mathcal{A}_n \cup \{\nabla_0\} = \{\nabla_0, \nabla_1, \dots, \nabla_n\}$:

$$\begin{array}{ll}
\mathcal{B}_n\ (n{=}2m): & \nabla_0 : \mathrm{A}_0 \to \mathrm{A}_0 \\
& \nabla_1 : \mathrm{A}_0 \to \mathrm{A}_1 \\
& \nabla_2 : \mathrm{A}_1 \to \mathrm{A}_2 \\
& \vdots \\
& \nabla_i : \mathrm{A}_{i-1} \to \mathrm{A}_i \\
& \vdots \\
& \nabla_m : \mathrm{A}_{m-1} \to \mathrm{A}_m \\
& \nabla_{m+1} : \mathrm{A}_m \to \mathrm{A}_{m-1} \\
& \vdots \\
& \nabla_{n-j} : \mathrm{A}_{j+1} \to \mathrm{A}_j \\
& \vdots \\
& \nabla_{n-1} : \mathrm{A}_2 \to \mathrm{A}_1 \\
& \nabla_n : \mathrm{A}_1 \to \mathrm{A}_0,
\end{array}
\qquad
\begin{array}{ll}
\mathcal{B}_n\ (n{=}2m{+}1): & \nabla_0 : \mathrm{A}_0 \to \mathrm{A}_0 \\
& \nabla_1 : \mathrm{A}_0 \to \mathrm{A}_1 \\
& \nabla_2 : \mathrm{A}_1 \to \mathrm{A}_2 \\
& \vdots \\
& \nabla_i : \mathrm{A}_{i-1} \to \mathrm{A}_i \\
& \vdots \\
& \nabla_m : \mathrm{A}_{m-1} \to \mathrm{A}_m \\
& \nabla_{m+1} : \mathrm{A}_m \to \mathrm{A}_m \\
& \nabla_{m+2} : \mathrm{A}_m \to \mathrm{A}_{m-1} \\
& \vdots \\
& \nabla_{n-j} : \mathrm{A}_{j+1} \to \mathrm{A}_j \\
& \vdots \\
& \nabla_{n-1} : \mathrm{A}_2 \to \mathrm{A}_1 \\
& \nabla_n : \mathrm{A}_1 \to \mathrm{A}_0.
\end{array}
\tag{20}$$

Let us define *higher order differential operations with Gateaux derivative* as the meaningful compositions of higher order of the functions from the set $\mathcal{B}_n = \{\nabla_0, \nabla_1, \dots, \nabla_n\}$. Our aim is to determine the number of higher order differential operations with Gateaux derivative. Let us define a binary relation σ "*to be in composition*":

$$\nabla_i \, \sigma \, \nabla_j = \begin{cases} \top\ , & (i{=}0 \wedge j{=}0) \vee (i{=}n \wedge j{=}0) \vee (j{=}i{+}1) \vee (i{+}j{=}n{+}1); \\ \bot\ , & \text{otherwise.} \end{cases} \tag{21}$$

and let $\mathtt{B} = [b_{ij}] \in \{0, 1\}^{(n+1)\times n}$ be the adjacency matrix associated with the graph which is determined by relation σ. So, analogously to the paper [6], the following statements hold.

Theorem 4.1. *Let $Q_n(\lambda) = |\mathtt{B} - \lambda \mathtt{I}| = \beta_0 \lambda^{n+1} + \beta_1 \lambda^n + \cdots + \beta_{n+1}$ be the characteristic polynomial of the matrix $\mathtt{B}$ and $v_{n+1} = [\,1 \ \dots \ 1\,]_{1\times(n+1)}$. If $\mathtt{g}(k)$ is the number of the kth order differential operations with Gateaux derivative, then the following formulas hold:*

$$\mathtt{g}(k) = v_{n+1} \cdot \mathtt{B}^{k-1} \cdot v_{n+1}^T \tag{22}$$

and

$$\beta_0 \mathtt{g}(k) + \beta_1 \mathtt{g}(k-1) + \cdots + \beta_{n+1} \mathtt{g}(k-(n+1)) = 0 \quad (k > n{+}1). \tag{23}$$

Lemma 4.2. *Let $Q_n(\lambda)$ and $P_n(\lambda)$ be the characteristic polynomials of the matrices $\mathtt{B}$ and $\mathtt{A}$ respectively. Then the following equality holds:*

$$Q_n(\lambda) = \lambda^2 P_{n-2}(\lambda) - \lambda P_n(\lambda). \tag{24}$$

Proof. Let us calculate the characteristic polynomial

$$Q_n(\lambda) = |\mathtt{B} - \lambda \mathtt{I}| = \begin{vmatrix} 1-\lambda & 1 & 0 & 0 & \dots & 0 & 0 & 0 & 0 \\ 0 & -\lambda & 1 & 0 & \dots & 0 & 0 & 0 & 1 \\ 0 & 0 & -\lambda & 1 & \dots & 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 & \dots & 0 & -\lambda & 1 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & 0 & -\lambda & 1 \\ 1 & 1 & 0 & 0 & \dots & 0 & 0 & 0 & -\lambda \end{vmatrix}. \tag{25}$$

Expanding the determinant $Q_n(\lambda)$ by the first column we have

$$Q_n(\lambda) = (1-\lambda)P_n(\lambda) + (-1)^{n+2}D_n(\lambda), \tag{26}$$

where

$$D_n(\lambda) = \begin{vmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & 0 \\ -\lambda & 1 & 0 & 0 & \dots & 0 & 0 & 0 & 1 \\ 0 & -\lambda & 1 & 0 & \dots & 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 & \dots & -\lambda & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & -\lambda & 1 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 & -\lambda & 1 \end{vmatrix}. \tag{27}$$

Let us expand the determinant $D_n(\lambda)$ by the first row and then in the next step, multiply the first row by -1 and add it to the last row. We obtain the determinant of order $n-1$:

$$D_n(\lambda) = \begin{vmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & 1 \\ -\lambda & 1 & 0 & 0 & \dots & 0 & 0 & 1 & 0 \\ 0 & -\lambda & 1 & 0 & \dots & 0 & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 & \dots & -\lambda & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & -\lambda & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 & -\lambda & 0 \end{vmatrix}. \tag{28}$$

Expanding the previous determinant by the last column we have

$$D_n(\lambda) = (-1)^n \begin{vmatrix} -\lambda & 1 & 0 & 0 & \dots & 0 & 0 & 0 & 1 \\ 0 & -\lambda & 1 & 0 & \dots & 0 & 0 & 1 & 0 \\ 0 & 0 & -\lambda & 1 & \dots & 0 & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 & \dots & 0 & -\lambda & 1 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 & -\lambda & 1 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & -\lambda \end{vmatrix}. \tag{29}$$

If we expand the previous determinant by the last row and if we expand the obtained determinant by the first column, we have the determinant of order $n-4$:

$$D_n(\lambda) = (-1)^n \lambda^2 \begin{vmatrix} -\lambda & 1 & 0 & 0 & \dots & 0 & 0 & 0 & 1 \\ 0 & -\lambda & 1 & 0 & \dots & 0 & 0 & 1 & 0 \\ 0 & 0 & -\lambda & 1 & \dots & 0 & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 & \dots & 0 & -\lambda & 1 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 & -\lambda & 1 \\ 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & -\lambda \end{vmatrix}. \tag{30}$$

In other words

$$D_n(\lambda) = (-1)^n \lambda^2 P_{n-4}(\lambda). \tag{31}$$

From equalities (31) and (26) there follows:

$$Q_n(\lambda) = (1-\lambda)P_n(\lambda) + \lambda^2 P_{n-4}(\lambda). \tag{32}$$

On the basis of Lemma 3.2. the following equality holds:

$$Q_n(\lambda) = \lambda^2 P_{n-2}(\lambda) - \lambda P_n(\lambda). \tag{33}$$

□

Lemma 4.3. *Let $Q_n(\lambda)$ be the characteristic polynomial of the matrix* B. *Then the following recurrence holds:*

$$Q_n(\lambda) = \lambda^2 \big(Q_{n-2}(\lambda) - Q_{n-4}(\lambda)\big). \tag{34}$$

Proof. On the basis of Lemma 3.2. and Lemma 4.2. the Lemma follows. □

Lemma 4.4. *Let $Q_n(\lambda)$ be the characteristic polynomial of the matrix* B. *Then it has the following explicit form*:

$$Q_n(\lambda) = \begin{cases} (\lambda-2) \displaystyle\sum_{k=1}^{\lfloor \frac{n}{4} \rfloor + 1} (-1)^{k-1} \binom{\frac{n+1}{2} - k}{k-1} \lambda^{n-2k+2} & , \ n = 2m+1; \\ \displaystyle\sum_{k=1}^{\lfloor \frac{n+3}{4} \rfloor + 2} (-1)^{k-1} \left(\binom{\frac{n}{2} - k + 2}{k-1} + \binom{\frac{n}{2} - k + 2}{k-2} \lambda \right) \lambda^{n-2k+3} & , \ n = 2m. \end{cases} \tag{35}$$

Proof. On the basis of Lemma 3.3 and Lemma 4.2. the Lemma follows. □

The recurrences for dimensions $n = 3, 4, \dots, 10$ are obtained by means of Malešević-Jovović [7] and they are given in the table below.

Dimension	Recurrence for the num. of the kth order diff. operations with Gateaux derivative
$n = 3$	$\mathsf{g}(k) = 2\mathsf{g}(k-1)$
$n = 4$	$\mathsf{g}(k) = \mathsf{g}(k-1) + 2\mathsf{g}(k-2) - \mathsf{g}(k-3)$
$n = 5$	$\mathsf{g}(k) = 2\mathsf{g}(k-1) + \mathsf{g}(k-2) - 2\mathsf{g}(k-3)$
$n = 6$	$\mathsf{g}(k) = \mathsf{g}(k-1) + 3\mathsf{g}(k-2) - 2\mathsf{g}(k-3) - \mathsf{g}(k-4)$
$n = 7$	$\mathsf{g}(k) = 2\mathsf{g}(k-1) + 2\mathsf{g}(k-2) - 4\mathsf{g}(k-3)$
$n = 8$	$\mathsf{g}(k) = \mathsf{g}(k-1) + 4\mathsf{g}(k-2) - 3\mathsf{g}(k-3) - 3\mathsf{g}(k-4) + \mathsf{g}(k-5)$
$n = 9$	$\mathsf{g}(k) = 2\mathsf{g}(k-1) + 3\mathsf{g}(k-2) - 6\mathsf{g}(k-3) - \mathsf{g}(k-4) + 2\mathsf{g}(k-5)$
$n = 10$	$\mathsf{g}(k) = \mathsf{g}(k-1) + 5\mathsf{g}(k-2) - 4\mathsf{g}(k-3) - 6\mathsf{g}(k-4) + 3\mathsf{g}(k-5) + \mathsf{g}(k-6)$

The values of the function $\mathsf{g}(k)$, for small values of the argument k, are given in the database of integer sequences [8] as the following sequences A000079 ($n = 3$), A090990 ($n = 4$), A007283 ($n = 5$), A090992 ($n = 6$), A000079 ($n = 7$), A090994 ($n = 8$), A020714 ($n = 9$), A129638 ($n = 10$).

References

[1] Ivanov, A. B., Vector analysis, in M. Hazewinkel, ed., *Encyclopaedia of Mathematics*, Springer, 2002. Text available at http://eom.springer.de/V/v096360.htm.

[2] G. A. Korn and T. M. Korn, *Mathematical Handbook for Scientists and Engineers: Definitions, Theorems, and Formulas for Reference and Review*, Courier Dover Publications, 2000.

[3] B. J. Malešević, A note on higher-order differential operations, *Univ. Beograd, Publ. Elektrotehn. Fak., Ser. Mat.* **7** (1996), 105–109. Text available at http://pefmath2.etf.bg.ac.yu/files/116/846.pdf.

[4] B. J. Malešević, Some combinatorial aspects of differential operation composition on the space R^n, *Univ. Beograd, Publ. Elektrotehn. Fak., Ser. Mat.* **9** (1998), 29–33. Text available at http://pefmath2.etf.bg.ac.yu/files/118/869.pdf.

[5] S. Basov, *Multidimensional Screening*, Springer 2005.

[6] B. J. Malešević, Some combinatorial aspects of the composition of a set of function, *Novi Sad J. Math.*, **36** (1), 2006, 3–9. Text available at http://www.im.ns.ac.yu/NSJOM/Papers/36_1/NSJOM_36_1_003_009.pdf.

[7] B. J. Malešević and I. V. Jovović, *A procedure for finding the kth power of a matrix.* http://www.maplesoft.com/Applications.

[8] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences.* http://www.research.att.com/~njas/sequences/.

2000 *Mathematics Subject Classification*: 05C30, 26B12, 58C20.

Keywords: compositions of the differential operations, enumeration of graphs and maps, Gateaux directional derivative.

(Concerned with sequences A000079, A007283, A020701, A020714, A090989, A090990, A090991, A090992, A090993, A090994, A090995, and A129638.)

Received June 5 2007; revised version received July 30 2007. Published in *Journal of Integer Sequences*, August 3 2007.

Return to Journal of Integer Sequences home page.