Journal of Integer Sequences, Vol. 15 (2012),
Article 12.4.3

A Generalized Apéry Series

M. L. Glasser
Department of Physics
Clarkson University
Postdam, NY 13699-5820
USA
laryg@clarkson.edu

Abstract

The inverse central binomial series

$$S_k(z) = \sum_{n=1}^{\infty} \frac{n^k z^n}{\binom{2n}{n}},$$

popularized by Apéry and Lehmer, is evaluated for positive integers k along with the asymptotic behavior for large k. We show that the value $z = 2$, as commented on by D. H. Lehmer, provides a unique relation to π.

1 Introduction

Since the appearance of $S_{-3}(1)$ in Apéry's famous proof [1] in 1979 that $\zeta(3)$ is irrational, an extensive literature has been devoted to the series

$$S_k(z) = \sum_{n=1}^{\infty} \frac{n^k z^n}{\binom{2n}{n}} \tag{1}$$

For example, in 1985 Lehmer [2] presented a number of special cases which could be obtained from the Taylor series for $f(x) = x^{-1/2}(1-x)^{-1/2}\sin^{-1} x$ using only elementary calculus. In passing, he noted that when k is a positive integer, $S_k(2)$ had the form $a_k - b_k\pi$ and that the rational number a_k/b_k "is a close approximation to π. This remark was recently taken up by Dyson et al. [3], who proved that $|a_k/b_k - \pi| = O(Q^{-k})$ as $k \to \infty$ where $Q = \sqrt{1 + (2\pi/\ln 2)^2}$. Lehmer also showed that for positive integer k

$$S_k(z) = \frac{2^{k+z^{5/2}} z^{1/2}}{(4-z)^{k+3/2}}(A_k(z/4)\sin^{-1}(\sqrt{z/4}) + \sqrt{z(4-z)}B_k(z/4)) \tag{2}$$

where A_k and B_k are recursively defined polynomials. It was apparently not until 2005 that (2) was evaluated explicitly, for $z = 1$, by J. Borwein and P. Girgensohn [4] who showed

$$S_k(1) = \frac{1}{2}(-1)^{k+1}\sum_{j=1}^{k+1}(-1)^j j! S(j+1,j)3^{-j}\binom{2j}{j}\left(\sum_{i=1}^{j-1}\frac{3^i}{(2i+1)\binom{2i}{i}} + \frac{2}{3\sqrt{3}}\pi\right). \quad (3)$$

where the *Stirling numbers of the second kind* are defined by

$$S(k,j) = \frac{(-1)^j}{j!}\sum_{m=0}^{j}(-1)^m m^k\binom{j}{m}. \quad (4)$$

The aim of the present note is to extend (3) to complex z and thus to continue (1) analytically beyond its circle of convergence $|z| = 4$.

2 Calculation

We begin with the observation that $\left(m\binom{2m}{m}\right)^{-1} = B(m, m+1)$, where B denotes Euler's beta integral. Hence,

$$S_k(z) = \int_0^1 \frac{dt}{t}\sum_{m=1}^{\infty} m^{k+1}(zt(1-t))^m. \quad (5)$$

Next, equation (21) of Girgensohn and Borwein [4],

$$\sum_{m=1}^{\infty} m^p X^m = \sum_{n=1}^{p}\sum_{m=1}^{n}(-1)^{m+n}\binom{n}{m}m^p X^n(1-X)^{-n-1}, \quad (6)$$

gives

$$S_k(z) = \sum_{n=1}^{k+1}\sum_{m=1}^{n}(-1)^{m+n}\binom{n}{m}m^{k+1}\int_0^1 \frac{dt}{t}\frac{(zt(1-t))^n}{(1-zt(1-t))^{n+1}}. \quad (7)$$

In the appendix it is shown that

$$\int_0^1 \frac{dt}{t}\frac{(zt(1-t))^n}{(1-zt(1-t))^{n+1}} = \frac{\sqrt{\pi}\Gamma(n)}{\Gamma(n+1/2)}X^n \, {}_2F_1(-1/2, n; n+1/2; -X) \quad (8)$$

where $X = z/(4-z)$, so

$$S_k(z) = \sum_{n=1}^{k+1} n! B(n, 1/2) S(k+1, n) X^n \, {}_2F_1(-1/2, n; n+1/2; -X). \quad (9)$$

By induction, starting with the tabulated value for $n = 1$ and using Gauss' contiguity relations we find (some details are given in the appendix)

$${}_2F_1(-1/2, n; n+1/2; -X) =$$

$$\left(\frac{1}{2}\right)_n \left(\frac{1}{n!} + \frac{1}{\sqrt{\pi}\Gamma(n)} \sum_{k=0}^{n-1} \frac{(-1)^k \Gamma(k+1/2)}{(k+1)!} \binom{n-1}{k} \left(\frac{X+1}{X}\right)^{k+1} \times \right.$$
$$\left. \left[\sqrt{X} \sin^{-1} \sqrt{\frac{X}{X+1}} - \frac{1}{2} \sum_{l=1}^{k} \frac{(l-1)!}{(1/2)_l} \left(\frac{X}{X+1}\right)^l \right]\right). \tag{10}$$

(We have used the ascending factorial notation $(a)_n = \Gamma(a+n)/\Gamma(a)$). Therefore we have the principal result

$$S_k(z) = \sum_{n=1}^{k+1} n! \left(\frac{z}{4-z}\right)^n S(k+1,n) \times$$

$$\left(\frac{1}{n} + \sum_{p=0}^{n-1} (-1)^p \frac{(1/2)_p}{(p+1)!} \binom{n-1}{p} \left(\frac{4}{z}\right)^{p+1} \left(\sqrt{\frac{z}{4-z}} \sin^{-1} \frac{\sqrt{z}}{2} - \frac{1}{2} \sum_{l=1}^{p} \frac{\Gamma(l)}{(1/2)_l} \left(\frac{z}{4}\right)^l \right)\right) \tag{11}$$

Equation (11) is rather condensed; in unpacking it, sums with upper limit less than the lower limit are to be interpreted as 0. It is clear from (11) that for *rational* z

$$\sum_{m=1}^{\infty} \frac{n^k z^n}{\binom{2n}{n}} = R_1(z,k) + R_2(z,k) \sqrt{\frac{z}{4-z}} \sin^{-1} \frac{\sqrt{z}}{2}, \tag{12}$$

where R_j is a rational number.

One sees from (11) that $S_k(z)$ is analytic on the two-sheeted Riemann surface formed by two planes cut and rejoined along the real half-line $x > 4$. The numbers in (12) have the explicit expressions

$$R_1(z,k) = \tag{13}$$

$$\sum_{n=1}^{k+1} n! S(k+1,n) \left(\frac{z}{4-z}\right)^n \left(\frac{1}{n} - \frac{1}{2} \sum_{p=1}^{n-1} \sum_{l=1}^{p} \frac{(-1)^p (1/2)_p}{(p+1)!(1/2)_l} \binom{n-1}{p} \Gamma(l) \left(\frac{4}{z}\right)^{p-l+1}\right),$$

$$R_2(z,k) = \sum_{n=1}^{k+1} n! S(k+1,n) \sum_{p=0}^{n-1} \frac{(-1)^p}{(p+1)!} \binom{n-1}{p} \left(\frac{4}{z}\right)^{p+1}. \tag{14}$$

3 Asymptotics

It is convenient to work in terms of the exponential generating function

$$G(z,t) := \sum_{k=0}^{\infty} S_k(z) \frac{t^k}{k!} = S_0(ze^t) = \frac{z}{4-ze^t} + \frac{4\sqrt{z}e^{t/2}}{(4-ze^t)^{3/2}} \sin^{-1} \frac{\sqrt{z}e^{t/2}}{2} \tag{15}$$

To find the generating functions $\rho_j(z,t) := \sum R_j(z,k) t^k/k!$, it would be simplest to start with a series $D_k(z) = R_1(z,k) - R_2(z,k) \sqrt{\frac{z}{4-z}} \sin^{-1} \sqrt{z}/2$, work out its generating function

$D(z,t)$ and by taking the sum and difference identify ρ_1 and ρ_2. However, this series has not been found and there is nothing to guarantee its existence in tractable form. Therefore, the ρ_j were evaluated directly from (13) and (14). The details are omitted as the results

$$\begin{aligned}\rho_1(z,t) &= \frac{ze^t}{4-ze^t} + \\ &\frac{8}{\pi}\sqrt{\frac{ze^t}{(4-ze^t)^{3/2}}}\left(\sin^{-1}\frac{\sqrt{z}e^{t/2}}{2}\cos^{-1}\frac{\sqrt{z}}{2} - \cos^{-1}\frac{\sqrt{z}e^{t/2}}{2}\sin^{-1}\frac{\sqrt{z}}{2}\right), \quad (16)\end{aligned}$$

$$\rho_2(z,t) = 4\sqrt{\frac{(4-z)e^t}{(4-ze^t)^3}} \quad (17)$$

are easily verified. In the case $z=2$, (15) and (16) are identical to Dyson's formulas [3, 5] obtained empirically.

In view of the prominent role that the ratio $R_1(z,t)/R_2(z,t)$ plays in Dyson et al. [3] for $z=2$ it is interesting to examine it for general z. From (17) we have

$$R_2(z,k) = \frac{2k!\sqrt{4-z}}{\pi i}\oint\frac{ds}{s^{k+1}}\frac{e^{s/2}}{(4-ze^s)^{3/2}}. \quad (18)$$

The non-zero singularity closest to $s=0$ is $s_0=\ln(4/z)$ and it dominates the asymptotic behavior. Ignoring the other singularities, distorting the contour to a small circle about s_0 and translating back to the origin by $t=s-s_0$, we have

$$R_2(z,k) \sim -\frac{k!\sqrt{4-z}}{zs_0^{k+1}}\oint\frac{dt}{2\pi i}\frac{e^{t/2}}{(1-e^t))^{3/2}}. \quad (19)$$

The exact value of the integral in (19) is $-(2/\pi)\sqrt{e/(e-1)}$, and so

$$R_2(z,k) \sim \frac{k!}{(\ln(4/z))^{k+1}}\frac{2}{\pi}\sqrt{\frac{e(4-z)}{z(e-1)}}. \quad (20)$$

In the same way we obtain

$$R_1(z,k) \sim \frac{k!}{(\ln(4/z))^{k+1}}\left(\sqrt{2}+\frac{2}{\pi}\left(\sqrt{\frac{e}{e-1}}-\sqrt{2}\right)\cos^{-1}\frac{\sqrt{z}}{2} - \frac{2^{3/2}}{\pi}\sin^{-1}\frac{\sqrt{z}}{2}\right). \quad (21)$$

4 Discussion

From (20) and (21) we find

$$\lim_{k\to\infty}\left(\frac{R_1(z,k)}{R_2(z,k)} - \sqrt{\frac{z}{4-z}}\sin^{-1}\frac{\sqrt{z}}{2}\right) = \sqrt{\frac{z}{4-z}}\left(\cos^{-1}\frac{\sqrt{z}}{2} - \sin^{-1}\frac{\sqrt{z}}{2}\right). \quad (22)$$

It thus appears that Lehmer's choice, $z = 2$, is the unique permissible case for which the limit (22) vanishes. (Also the *Lehmer limit*, as defined by Dyson et al. [3], relates to $\pi/4$ here rather than π). Finally, for negative integer indices, since

$$2S_{-k}(z) = {}_{k+1}F_k(1, \ldots, 1; \tfrac{3}{2}, 2, \ldots, 2; \tfrac{1}{4}z), \tag{23}$$

the fact that $S_{-k}(z)$ can be obtained from $S_0(z)$ by successive integrations with respect to z and the explicit evaluations by Lehmer [2], Borwein and Girgensohn [4] and others [6, 7, 8, 9, 10] it should be possible to obtain explicit values for sundry generalized hypergeometric functions.

5 Appendix: Derivation of Equations (8) and (10)

Let us consider, for any integrable function F,

$$I = \int_0^1 \frac{dt}{t} F(t(1-t))$$

Let $u = t(1-t)$, so $u(0) = u(1) = 0$; $u(1/2) = 1/4$. Then there are two expressions for t:

$$t_+ = \frac{1}{2}(1 + \sqrt{1-4t}) \quad \text{for } \frac{1}{2} \le t \le 1, \text{ with } \frac{dt_+}{t_+} = \left(1 - \frac{1}{\sqrt{1-4u}}\right)\frac{du}{u}$$

and

$$t_- = \frac{1}{2}(1 - \sqrt{1-4t}) \quad \text{for } 0 \le t \le \frac{1}{2}, \text{ with } \frac{dt_+}{t_+} = \left(1 + \frac{1}{\sqrt{1-4u}}\right)\frac{du}{u}.$$

Consequently,

$$\begin{aligned} I &= \int_0^{1/2} \frac{dt_-}{t_-} F(u) + \int_{1/2}^1 \frac{dt_+}{t_+} F(u) = 2\int_0^{1/4} \frac{du}{u\sqrt{1-4u}} F(u) \\ &= 2\int_0^1 \frac{dx}{x\sqrt{1-x}} F\left(\frac{1}{4}x\right) = 2\int_0^1 \frac{dt}{(1-t)\sqrt{t}} F\left(\frac{1-t}{4}\right) \end{aligned}$$

and, with $t = x^2$,

$$I = 4\int_0^1 \frac{dx}{1-x^2} F\left(\frac{1-x^2}{4}\right).$$

Therefore,

$$L = \int_0^1 \frac{dt}{t} \frac{(zt(1-t))^\alpha}{(1-zt(1-t))^\beta} = 2\left(\frac{z}{4}\right)^{\alpha-\beta} \int_0^1 dx \frac{(1-x^2)^{\alpha-1}}{(a^2+x^2)^\beta},$$

where $a^2 = 1/X = (4-z)/z$.

From standard references

$$\int_0^1 dx \cos(xy)(1-x^2)^{\alpha-1} = \sqrt{\frac{\pi y}{8}} \left(\frac{2}{y}\right)^\alpha \Gamma(\alpha) J_{\alpha-1/2}(y),$$

$$\int_0^\infty dx \cos(xy)(a^2+x^2)^{-\beta} = \frac{\sqrt{\pi}}{\Gamma(\beta)}\left(\frac{y}{2a}\right)^{\beta-1/2} K_{\beta-1/2}(ay)$$

so, by the Parseval relation for the Fourier transform

$$\int_0^1 dx \frac{(1-x^2)^{\alpha-1}}{(a^2+x^2)^\beta} = \frac{2^{\alpha-\beta}}{a^{\beta-1/2}}\frac{\Gamma(\alpha)}{\Gamma(\beta)}\int_0^\infty dy y^{\beta-\alpha} J_{\alpha-1/2}(y) K_{\beta-1/2}(ay).$$

This is a tabulated Hankel Transform and yields

$$L = \sqrt{\pi}\left(\frac{4}{z}\right)^{\beta-\alpha} X^\beta \frac{\Gamma(\alpha)}{\Gamma(\alpha+1)}\ {}_2F_1\left(\frac{1}{2},\beta;\alpha+\frac{1}{2};-X\right).$$

Consequently

$$\int_0^1 \frac{dx}{x}\frac{(zt(1-t))^n}{(1-zt(1-t))^{n+1}} = \sqrt{\pi}\left(\frac{4}{z}\right) X^{n+1}\ {}_2F_1\left(\frac{1}{2},n+1;n+\frac{1}{2};-X\right)$$

However, since ${}_2F_1(a,b;c;z) = (1-z)^{c-a-b}\ {}_2F_1(c-a,c-b;c;z)$,

$${}_2F_1\left(\tfrac{1}{2},n+1;n+\tfrac{1}{2};-X\right) = (1+X)^{-1}\ {}_2F_1\left(-\tfrac{1}{2};n;n+\tfrac{1}{2};-X\right)$$

Next, we note that [11, p . 590]

$${}_2F_1(-1/2,1;3/2;z) = \frac{1}{2}\left(1+(1-z)\frac{\tanh^{-1}\sqrt{z}}{\sqrt{z}}\right).$$

With $z \to -z$, noting that $-i\tanh^{-1} iw = \sin^{-1}\sqrt{\frac{w}{1+w}}$ one has

$${}_2F_1(-1/2,1;3/2;-z) = \frac{1}{2}(1+(1+z)\frac{\sin^{-1}\sqrt{\frac{z}{1+z}}}{\sqrt{z}}). \tag{24}$$

We next apply Gauss' differentiation formula

$$\frac{d}{dz}((1+z)^k\ {}_2F_1(-1/2,k;k+1/2;-z)) =$$

$$\frac{2k(k+1)}{2k+1}(1+z)\ {}_2F_1(-1/2,k+1;k+3/2;-z). \tag{25}$$

Iteration of (25) starting with (24), after a great deal of tedious algebra, aided by Mathematica, results in (10).

6 Acknowledgements

The author is grateful to Profs. N. E. Frankel for suggesting this problem and F. J. Dyson for enlightening correspondence.

References

[1] R. Apéry, Irrationalité de $\zeta(2)$ et $\zeta(3)$, *Asterique* **61** (1979), 11–13.

[2] D. H. Lehmer, Interesting series involving the central binomial coefficient, *Amer. Math. Monthly* **92** (1985), 449–457.

[3] F. J. Dyson, N. E. Frankel, and M. L. Glasser, Lehmer's second interesting series, preprint. Available at http//:arxiv.org/abs/1009.4274.

[4] J. M. Borwein and R. Girgensohn, Evaluations of binomial series, *Aequ. Math.* **70** (2005), 25–36.

[5] F. J. Dyson, personal correspondence.

[6] I. J. Zucker, On the series $\sum \binom{2n}{n}^{-1} n^{-k}$ and related series, *J. Number Theory* **20** (1981), 92–102.

[7] N. Batir, On the series $\sum \binom{2k}{k}^{-1} k^{-a} x^k$, *Proc. Indian Acad. Sci.* **115** (2005), 371–382.

[8] J. -H. Yang and F. -Z. Zhao, Certain sums involving inverses of binomial coefficients and some integrals, *J. Integer Sequences* **10** (2007), 1–11.

[9] R. Sprugnoli, Sums of the reciprocals of the central binomial coefficients, *INTEGERS* **6** (2006), 1–18.

[10] R. J. Mathar, Corrigenda to "Interesting series involving the central binomial coefficient by D. H. Lehmer, preprint. Available at http//:arxiv.org/abs/0905.0215v1.

[11] A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, *Tables of Integrals and Series* Vol. 3, Gordon and Breach, 1998.

2000 *Mathematics Subject Classification*: Primary 11B65; Secondary 33B05.
Keywords: Binomial coefficient, infinite series, generating function.

(Concerned with sequences A008277 and A145557.)

Received October 8 2011; revised version received April 2 2012. Published in *Journal of Integer Sequences*, April 6 2012.

Return to Journal of Integer Sequences home page.